

# Air Quality Governance in the ENPI East Countries

## *RPP3 – Training workshop*

### Spatial distribution

14-16 July, 2014, Chişinău, Moldova



MWH



# Outline

- Introduction
- Procedure for spatial distribution of emissions
- Input data
  - Emissions
  - Spatial data
- Examples of spatial data
  - Best case
  - What we got
- Output
  - Emission maps
  - Input to air quality modelling

# Spatial distribution of emissions

## Introduction (1)

- To add a spatial component to the emission inventories, it is necessary to include various digitized spatial data
- Depending on the availability of spatial data, separate spatial distributions can be made for:
  - a single source
  - a group of sources with comparable properties
  - an emission sector
- Spatial emissions must be consistent with the emission inventory for the city

# Spatial distribution of emissions

## Introduction (2)

- The spatial data have to be in a format compatible with a Geographical Information System (GIS)
  - GIS
    - ArcGIS or Quantum GIS (open source)
  - Data format
    - Shape files
    - personal geodatabase
    - file geodatabase
    - raster files
  - Datum and projection must be specified for all data sets

# Spatial distribution of emissions Procedure

- Definition of city boundaries
  - City, municipality, etc.
  - Correspond to the available statistical data, if possible
- Identification of point sources within the boundaries
  - Power plants, industrial plants, refineries, etc.
- Identification of area sources within the boundaries
  - Small scale combustion, road traffic, etc.



# Input data

## Point sources

- The spatial model for point sources uses the coordinates for the specific plant and the calculated emissions
- Geographical coordinates can be derived from address using Google Earth
  - Longitude / Latitude (decimal degrees)
  - Datum: WGS 84
- Example of emission model input from point sources to the spatial distribution

SNAP	Long	Lat	Datum	Stack height	NO <sub>x</sub>	SO <sub>2</sub>	NMVOC	CO	TSP	PM <sub>10</sub>	PM <sub>2.5</sub>
03	<i>DD.dddd</i>	<i>DD.dddd</i>	WGS84	100							
03	<i>DD.dddd</i>	<i>DD.dddd</i>	WGS84	100							
01	<i>DD.dddd</i>	<i>DD.dddd</i>	WGS84	150							
03	<i>DD.dddd</i>	<i>DD.dddd</i>	WGS84	100							

# Input data

## Area sources (1)

- The spatial model for area sources uses the calculated emissions for the specific sources and corresponding spatial distribution keys
- Example of emission model input from area sources to the spatial distribution




SNAP	NO <sub>x</sub>	SO <sub>2</sub>	NMVOC	CO	TSP	PM <sub>10</sub>	PM <sub>2.5</sub>
02							
03							
07							
0802							
0804							
0805							
0808							

# Input data

## Area sources (2)

- Area sources
  - Emissions that are spatially associated with an area (a polygon)
    - E.g. residential heating, non-road mobile sources.
- Line sources
  - Emission sources that exhibits a line type of geography
    - E.g. road, railway, pipeline or shipping lane
- Grids
  - Net of squares, in this case with a resolution of 1 km x 1 km
  - Point, line and polygon features can be combined to grids and then several different layers of information (emission sources) can be aggregated

# Combining different spatial features






# Input data

## Examples of spatial data (1)

- Road traffic
  - Road network
    - Road classification
      - Motorway / urban road / rural road
      - Motorway / major road / minor road
      - Street width categories
    - Mileage
      - Annual Average Daily Traffic
        - » per street segment
        - » per vehicle types

# Road Maps






# Input data

## Examples of spatial data (2)

- Small scale combustion
  - Land use map
    - Building types
      - Residential
      - Commercial / institutional
      - Industry
    - Heating districts
      - Share of buildings connected to district heating



# Land use Maps







# Input data

## What we got from cities

- Very different spatial data available from city to city
- Some spatial data are available but could not be provided to the project for free

Feature	Number of countries	Feature	Number of countries
<b>Demarcation of the area of the city</b>	3	<b>Buildings</b>	2
<b>Administrative units</b>	1	Building types (tall / one-storage)	-
Population density	-	Urban areas	-
<b>Road network</b>	2	Rural areas	-
Road types (urban/rural/highway)	-	<b>Settlement areas</b>	1
Road width	-	Commercial areas	-
Annual average mileage	-	<b>Industrial areas</b>	1
Annual average mileage per vehicle type	-	Construction areas	-
Annual average number of vehicles	-	Heating districts	-
Annual average number of vehicles per vehicle type	-	Airport areas	-
<b>Railway network</b>	1	Port facilities	-

# Input data from cities





# Input data

## What we got elsewhere

- Due to the very limited spatial data provided for some cities, it was necessary to find alternative spatial data
  - OpenStreetMap (<http://download.geofabrik.de/>)
    - Road network, buildings, land use, etc.
  - Population density
    - Landsat (<http://web.ornl.gov/sci/landscan/index.shtml>)
    - Center for International Earth Science Information Network (CIESIN), Columbia University (<http://sedac.ciesin.columbia.edu/data/collection/gpw-v3>)
- These datasets were used even if the level of detail is low, to illustrate the preferred methodology, and since it in most cases provides a better result, than just distribution emissions evenly across the entire city area.

# Input data

## What we got elsewhere



# Spatial distribution key (1)

- Use GIS to:
  1. Intersect industrial area and grid
  2. Calculate area of new polygons (calculate geometry)
  3. Calculate share of total industrial area for each polygon



**Industrial area**



**1 km x 1 km grid**



**Intersection of industrial area and 1 km x 1 km grid**

# Spatial distribution key (2)

- Example of spatial distribution keys based on road network, buildings and population density, respectively.

GridID	ShareOfLength
1km_6149_600	0,01691
1km_6150_599	0,00847
.....	.....
1km_6150_600	0,06
1km_6150_601	0,01
<b>Grand Total</b>	<b>1,00</b>

GridID	ShareOfBuildArea
1km_6151_598	0,00022
1km_6151_599	0,00045
.....	.....
1km_6151_607	0,028
1km_6151_608	0,034
<b>Grand Total</b>	<b>1,000</b>




GridID	ShareOfCapita
1km_6149_600	0,00114
1km_6149_601	0,00023
.....	.....
1km_6150_597	0,00564
1km_6150_599	0,00177
<b>Grand Total</b>	<b>1,00000</b>

# Output data

<b>TOTAL CITY EMISSIONS FOR AREA SOURCES</b>				
<b>SNAP</b>	<b>NOx</b>	<b>SO2</b>	<b>CO</b>	<b>TSP</b>
02	1403.229	7.016	701.614	51.452

<b>SPATIAL DISTRIBUTION KEY</b>		<b>GRIDDED EMISSIONS</b>			
<b>GridID</b>	<b>ShareOfArea</b>	<b>NOx</b>	<b>SO2</b>	<b>CO</b>	<b>TSP</b>
		<b>E(cell) = E(city) * ShareOfArea</b>			
1km_3737_511	0.0081	11.3927	0.0570	5.6964	0.4177
1km_3737_512	0.0005	0.6911	0.0035	0.3455	0.0253
1km_3738_458	0.0032	4.4288	0.0221	2.2144	0.1624
1km_3738_511	00032	4.4246	0.0221	2.2123	0.1622
1km_3738_512	0.0033	4.7005	0.0235	2.3503	0.1724
.....	.....	.....	.....	.....	.....
1km_3754_513	0.0000	0.0055	0.0000	0.0027	0.0002
1km_3754_514	0.0000	0.0596	0.0003	0.0298	0.0022
<b>SUM</b>	<b>1.0000</b>	<b>1403.229</b>	<b>7.016</b>	<b>701.614</b>	<b>51.452</b>

# Output Emission maps





Thank you for your attention



**MWH**



Center for Environmental and Water Research



**FORCE  
TECHNOLOGY**



WORLDWIDE CONSULTANTS



**NERI**  
AARHUS UNIVERSITY

